

FIRST OFF-SHORE SITE BIRD
MONITORING IN POLAND
(DEBKI-BIAŁOGÓRA, 2002–2004)

Przemysław Busse

ABSTRACT

Busse P. 2015. *First off-shore site bird monitoring in Poland (Dębki-Białogóra, 2002-2004)*. Ring 37: 19-54

Due to the presence along the Polish Baltic coast of migratory bird flyways from north-eastern European breeding grounds to wintering areas distributed on continental shelf waters of western and southwestern Europe, the area has been designated as a *NATURA 2000* site (PLB 990002). Therefore the site of a planned off-shore wind farm in this area requires monitoring as to its potential influence on birds. This was the first such monitoring performed in Poland. The expanse of water covered by the research included the planned location of the wind farm as well as adjacent areas. A series of 15 investigative cruises were undertaken from the beginning of October to the beginning of May. The period of observations was divided into five seasons: early autumn, late autumn, winter, early spring and late spring. The standard method of counting birds on transects in the form of strips reaching 300 m from the ship was used in the research, as well as the 'snapshot' technique (scan with bands transect with snapshot technique). Additional observations were made from a point on the shore.

In the study area maritime birds are present in low or moderate densities, with localized clusters. Two diving benthophagous species dominate: the Long-tailed Duck (58.9%) and the Velvet Scoter (34.3%). The next two commonest species have a share of over 1% within the community – the Common Scoter (3.2%) and the Herring Gull (2.4%). Other species are very uncommon. There is very high fluctuation in the number of birds both observed on the water surface and seen in the air. In the area studied no pronounced migratory passage of waterfowl was observed; the usual migratory flyways probably lie farther to the north and the birds observed in flight perform mainly local movements. Observed bird densities in the study area are considerably lower (52.0 ind./km²) than those estimated for the entire *NATURA 2000* area (116.7 ind./km²). In the area of the planned wind farm densities are even lower (36.1 ind./km², i.e. 31% of the *NATURA 2000* level), while densities in neighbouring areas are still below the *NATURA 2000* average.

In the subsequent administrative procedure the area was not accepted as the location of the planned wind farm.

P. Busse Bird Migration Research Foundation, Przebendowo 3, 84-210 Choczewo, Poland.
E-mail: busse@wbwp-fund.eu

Keywords: bird monitoring, off-shore, wind-farm, waterfowl, distribution, seasonal dynamics

INTRODUCTION

The Baltic Sea area is of great importance for several maritime bird species. These birds, breeding at high latitudes, migrate this way to moulting and wintering grounds located on the western and southern waters of the Baltic and the North Sea. The role of these areas has been described in a report addressed to the European Commission entitled '*Important Marine Areas for Wintering Birds in the Baltic Sea*' (Durinck *et al.* 1994). The '*Material*' chapter of this document provided data describing the basis for evaluation of different areas of the Baltic Sea for maritime birds. It can be seen here that the amount of original data collected in different parts of the area was highly varied, from single controls to detailed studies based on numerous observations made from the coast and from ships, as well as aerial censuses. The Polish coast was extremely under-studied – most evaluations were based on only one study cruise and only the Pomeranian Bay and Slupsk Bank were observed more than once. Therefore knowledge of the occurrence, distribution and migration of maritime birds in this area is insufficient for real evaluation of its importance for birds. Despite this, the entire area of Polish waters with water depths below 20 meters was designated as protected (OSO) under the NATURA 2000 project (PLB990002). This area is very large, covering 211,596 ha. The wintering population of birds was estimated at about 247,000 birds, which means an average density of about 117 ind./km².

Along the Baltic coasts there are migratory flyways from north-eastern European breeding grounds to wintering grounds distributed on the continental shelf waters of western and southwestern Europe. Migration – partly true migration and partly moult movements – begins as early as summer. As early as July one can observe the first migrating drakes of duck species that leave their breeding grounds after the females have started to breed. The most common duck species at this time is the Common Scoter, migrating to moult on shallow Danish waters. Knowledge of this migration is rather poor, but unpublished data show that it is not intensive on the southern Baltic waters. Migration of other waterfowl is very prolonged and lasts until the end of October or even the first half of November.

Spring migration of waterfowl is much quicker than in autumn and more intensive. The first migrating flocks can already be seen in the last days of February, and the peak of migration occurs in April. The latest migrants can be seen until the end of May, but are observed infrequently. Up to the time of the observations described here, research on spring migration on the Polish coast had been carried out only in 2000, near Rozewie (Meissner 2003). The number of migrating birds was rather high, on some days reaching as many as 1,500 individuals per hour. Ninety percent of all observed birds were gulls, and 80-90 percent of them flew within a 1 km strip from the shore. The only group of birds migrating at higher distances from land were divers (W. Meissner – unpublished data). These birds were observed mainly at distances of 500-1,500 meters from the beach. In this case the direction of flights was not parallel to the shoreline and the birds headed northeast from Rozewie. During 80 hours of observations 164 individuals were counted, including 34 Black-throated Divers, 64 Red-throated Divers and one Yellow-billed Diver (others were not identified). On

average 2.1 ind./hour passed the observation stand. In comparison with Estonian observations (178 ind./h – Kontkanen 1996), intensity of migration on the Polish coast is rather low.

The Polish maritime area had not been exploited for wind farms before the beginning of the 21st century and therefore no studies on birds were carried out to evaluate potential conflicts between this method of energy production and birds using the area. This indicates the need for special studies in this regard at planned sites for wind farms. The first application for permission to build a wind farm, near Dębki and Karwieńskie Błota, necessitated adequate studies at this location.

The evaluation below is based on observations performed by a team from ECOTONE Enterprise, Sopot, guided by Prof. Włodzimierz Meissner (Department of Ecology and Zoology of Vertebrates, Gdańsk University). The raw data are presented in reports by ECOTONE: *'Report from studies on numbers and distribution of birds observed at the area of planned wind farm Debki-Jastrzebia Góra'*, and *'Report from observations of migrating birds at Jastrzebia Góra-Dębki'*. Limited evaluation of these data was published earlier by Meissner (Meissner 2010). It was directed to three most numerous species only and covered nine of 15 cruises.

It must be stressed that the evaluation is a study of the local situation in this area and cannot be applied to other parts of the Polish Baltic coast. This is only a snapshot from the initial period of research on the distribution of ducks along the Polish Baltic coast. Since that time the numbers have changed and general estimations indicate much lower numbers of birds observed along the southern Baltic coast.

AREA AND METHODS

Observations of numbers and distributions of birds

Observations of the species composition, number and the distribution of waterfowl in the region of the planned investment were carried out on a rectangular expanse of water defined by the following coordinates:

- NW: 54°58'00"N 18°00'00"E
- NE: 54°58'00"N 18°18'00"E
- SW: 54°50'50"N 18°00'00"E
- SE: 54°50'50"N 18°18'00"E

The expanse of water covered by the research included the planned location of the wind farm as well as adjacent areas. Transects ran north-south along meridians 18°18', 18°12', 18°06' and 18°00' and were 13.5 km in length. Each transect was divided into 6 sections of about equal length, numbered from the shore towards the open sea. The planned location of the wind power stations was within the second sections and a small part of the thirds sections of the two central transects, as shown in Figure 1.

As previously decided, 15 investigative cruises were conducted, from the beginning of October to the beginning of May. The period of observations was divided into five seasons as shown in Table 1. To control for inter-year fluctuations, late autumn controls were done in both 2003 and 2004.




Fig. 1. The study area – summer observation point and ship cruises: vertical broken lines – transects, horizontal lines – observation sectors; ● – actual locations of wind turbines, ○ – line of wind turbines cancelled from building plans; isobaths of 10 and 20 m are shown as well as local depths.

Table 1
Phenological seasons and dates of exploration

Season	Dates of cruises
Early autumn	4 Oct. 2004
	25 Oct. 2004
Late autumn	4 Nov. 2004
	14 Nov. 2003
	28 Nov. 2003
	29 Nov. 2004
Winter	9 Dec. 2003
	30 Dec. 2003
	27 Jan. 2004
	27 Feb. 2004
Early spring	5 Mar. 2004
	28 Mar. 2004
	5 Apr. 2004
Late spring	20 Apr. 2004
	5 May 2004

The standard method of counting birds on transects in the form of strips reaching 300 m from the ship was used in the research, as well as the 'snapshot' technique (scan with bands transect with snapshot technique). This method is used nearly

worldwide in this type of research. A detailed description of the method is found in a manual by Komdeur *et al.* 1992.

The control covered an area of 16.2 km², or 0.675 km² per sector. Detail description of the area was published by Meissner (2010).

Observations of migration

The passage of waterfowl in the period from early autumn to late spring was observed during cruises aimed at estimating the density and distribution of birds (as above).

Summer observations were performed from a coastal stand on a 40 m cliff in Jas-trzębia Góra, near Rozewie. This location is suitable for such observations due to its high elevation and open view of the sea, which make it possible to follow birds flying a few kilometres from the shore (up to 2-2.5 km). This point is located outside the planned wind farm area. However, most birds, except for divers, migrate there parallel to the coast line, so they pass both the farm area and the observation point area. Observations were aimed in particular at waterfowl, but other large birds were counted as well. Birds were noted in three strips: up to 500 m, 500-1,000 m and more than 1,000 m from the coast. The estimation of a distance based on relations between a horizon distance in relation to elevation of an observer (Heinemann 1981). As the planned wind farm is to be about 4 km from the shore, the most important results are those obtained for the third strip. The birds were counted on 12 days, three times daily: in the morning (7.00-9.00), around midday (11.00-13.00) and in the afternoon (15.00-17.00). Observations were conducted continuously for at least one hour using a 20-60x80 telescope or binoculars. Altogether birds were counted for 41 hours and 45 minutes (Table 2).

Table 2
Time spent at the passage observation point at Rozewie in 2002. Hours:minutes

	Morning	Midday	Afternoon	Total
27 July	2:00	2:00	1:30	5:30
30 July	1:45	1:30	1:00	4:15
2 August	1:30	1:30	1:00	4:00
5 August	1:30	1:00	1:00	3:30
9 August	1:00	1:00	1:00	3:00
12 August	1:00	1:00	1:00	3:00
16 August	1:00	1:00	1:00	3:00
20 August	1:00	1:00	1:00	3:00
24 August	1:00	1:00	1:00	3:00
27 August	1:00	1:00	1:00	3:00
31 August	1:00	1:00	1:00	3:00
4 September	1:30	1:00	1:00	3:30
Total	15:15	14:00	12:30	41:45

Observations of migration of land birds were generally not conducted. Most land birds migrate over the sea during the night and at considerable heights. These obser-

vations require completely different techniques; moreover, migration normally takes place outside heights where collision with wind turbines may occur. Diurnal migration of land birds at heights comparable with those of windmill rotors could take place only rarely and in very special weather conditions, so the potential impact of the wind farm seems to be very low and cannot be properly estimated.

Analysis of data

Numbers of birds observed on the water surface are given here as densities per 1 km². The raw data were analysed in SURFER 8.0 computer software and presented as isolines describing the distribution of birds within the study area. Numbers for all bird species together, as well as for the two commonest species separately, were analysed in the program. The total picture of the distribution of birds for all seasons together was based on a weighted average of means for all seasons (weighted for length of the season: early autumn – 1 month, late autumn – 2 months, winter – 3 months, early spring – 1 month, and late spring – 1 month).

Intensity of flights is expressed as numbers of individuals observed flying per hour of observation, irrespective of whether they were observed from the shore or from a ship. This makes it possible to compare the results with literature data.

RESULTS

Numbers and distribution of birds

Altogether 11,116 maritime bird individuals were observed during the research cruises (Table 3). The two most common species – the Long-tailed Duck and Velvet Scoter – comprised 93.2% of all observed birds and attention was focused mainly on these species. Both species are diving benthophages, and the third most frequent species in the area, i.e. the Common Scoter (3.2% share in all maritime species), belongs to this ecological group as well. Diving ichthyophages (10 species) constitute less than 1% of birds observed in the area. Other species, apart from Herring Gull (1.2%), occur in negligible numbers.

Both total density and species densities fluctuate considerably between seasons (Table 4). The highest average density was found during early spring – it was about two times higher than in late autumn and winter and five times higher than in early autumn. This is a period of highly dynamic spring migration. Later in the year, during late spring, only small numbers of waterfowl were observed. However, Figure 2 shows that the average values may be not representative of the actual number dynamics, as control-to-control fluctuations in observed numbers are quite large. With only 15 controls conducted for all seasons together, a single observation of very high numbers could influence of the overall number dynamics and total densities. An example of this is the very high number of Long-tailed Duck observed on 14 November 2003 (119 ind./km², while the total average was only 29.1 ind./km²). This is confirmed by the high standard deviation ($SD = 31.9$) found for this species.

Table 3

Species composition and numbers of waterfowl in the area studied – total average.
+ – dominance less than 0.1%, density less than 0.1 ind./km²

	Total number	Dominance [%]	Density [n/km ²]
Diving benthophages			
Long-tailed Duck <i>Clangula hyemalis</i>	6551	58.9	28.4
Velvet Scoter <i>Melanitta fusca</i>	3817	34.3	16.5
Common Scoter <i>Melanitta nigra</i>	358	3.2	1.6
Common Eider <i>Somateria mollissima</i>	22	0.2	0.1
Diving ichthyophages			
Black-throated Diver <i>Gavia arctica</i>	2	+	+
Red-throated Diver <i>Gavia stellata</i>	2	+	+
Divers <i>indet. Gavia sp.</i>	1	+	+
Great Crested Grebe <i>Podiceps cristatus</i>	9	0.1	+
Red-necked Grebe <i>Podiceps grisegena</i>	1	+	+
Cormorant <i>Phalacrocorax carbo</i>	1	+	+
Goosanger <i>Mergus merganser</i>	7	0.1	+
Red-necked Merganser <i>Mergus serrator</i>	14	0.1	0.1
Razorbill <i>Alca torda</i>	29	0.3	0.1
Common Guillemot <i>Uria aalge</i>	4	+	+
Black Guillemot <i>Cephus grylle</i>	3	+	+
Others			
Great Black-backed Gull <i>Larus marinus</i>	15	0.1	0.1
Herring Gull <i>Larus argentatus</i>	268	2.4	1.2
Lesser Black-backed Gull <i>Larus fuscus</i>	2	+	+
Common Gull <i>Larus canus</i>	3	+	+
Black-headed Gull <i>Larus ridibundus</i>	1	+	+
Little Gull <i>Larus minutus</i>	2	+	+
Bean Goose <i>Anser fabalis</i>	3	+	+
Coot <i>Fulica atra</i>	1	+	+
Total	11 116	100	48.2

Apart from the overall density of birds and their number dynamics, the local distribution of the bird flocks is of utmost importance for evaluation of the area in terms of the location of the wind farm, as well as for studies on the feeding behaviour of the species observed. The local distribution of birds in the area is significantly uneven; when these distributions are tested using the chi-squared test, the probabilities obtained are undistinguishable from 0, e.g. for winter chi-squared ($df=23$) = 382, while the border value for $p = 0.01$ is only 41.6. It is generally accepted that densities of maritime birds, especially diving benthophages, depend on water depth. Verification of this statement on the data obtained, however, shows that this relation is not as clear and stable as might be expected. Table 5 shows that a negative correlation occurs between bird densities and the depth of the sea in autumn, but is absent in winter. This is true both for all bird species together and for the two most abundant

species. Therefore, the most important variable for birds seems to be concentration of food resources and not water depth alone. It is possible that more accessible resources in shallower waters are already exploited in autumn, while in winter the birds must use resources from greater depths. This hypothesis seems the most probable for the Long-tailed Duck.

Table 4
Densities of waterfowl in the study area in different seasons.
+ – density less than 0.1 ind./km²

	Seasons					Total
	Early autumn	Late autumn	Winter	Early spring	Late spring	
Diving benthophages						
Long-tailed Duck <i>Clangula hyemalis</i>	11.0	37.9	31.2	65.9	2.1	28.4
Velvet Scoter <i>Melanitta fusca</i>	8.1	20.4	20.4	32.7	3.0	16.5
Common Scoter <i>Melanitta nigra</i>	1.9	1.4	1.5	3.9	0.2	1.6
Common Eider <i>Somateria mollissima</i>	0.2	0.2		+		0.1
Density of diving benthophags	21.3	59.9	53.1	102.6	5.2	46.5
Diving ichthyophages						
Black-throated Diver <i>Gavia arctica</i>	+	+				+
Red-throated Diver <i>Gavia stellata</i>		+	+			+
Divers <i>indet. Gavia sp.</i>				+		+
Great Crested Grebe <i>Podiceps cristatus</i>	0.1	0.1	+			+
Red-necked Grebe <i>Podiceps grisegena</i>			+			+
Cormorant <i>Phalacrocorax carbo</i>	+					+
Goosanger <i>Mergus merganser</i>			0.1			+
Red-necked Merganser <i>Mergus serrator</i>		0.1	0.1			0.1
Razorbill <i>Alca torda</i>	0.2	0.3		0.1		0.1
Common Guillemot <i>Uria aalge</i>		0.1				+
Black Guillemot <i>Cepphus grylle</i>			+	+		+
Density of diving ichthyophags	0.3	0.6	0.4	0.2	0.0	0.3
Others						
Great Black-backed Gull <i>Larus marinus</i>	0.2	0.1	+			0.1
Herring Gull <i>Larus argentataus</i>	0.6	2.9	0.8	0.3	0.1	1.2
Lesser Black-backed Gull <i>Larus fuscus</i>					+	+
Common Gull <i>Larus canus</i>		+	+			+
Black-headed Gull <i>Larus ridibundus</i>				+		+
Little Gull <i>Larus minutus</i>					+	+
Bean Goose <i>Anser fabalis</i>					0.1	+
Coot <i>Fulica atra</i>					+	+
Density of others	0.9	3.0	0.9	0.3	0.3	1.3
Total density	22.5	63.4	54.4	103.1	5.5	48.2




Fig. 2. Densities of birds observed in each control – for all species together (TOTAL) and for the two most common species. Averages (m) are shown as a line and a 0.05 confidence interval ($\pm 2*SE$) as a dotted area.




Fig. 3. Distributions of all birds observed in the area during four successive controls in autumn 2004. Computer estimation (using SURFER 8.0) based on the data collected – isolines of densities expressed as number of individuals per km^2 .

Table 5

Correlation coefficients between sea depth and observed densities of birds. In bold – significant or highly significant correlations (< 0.05)

	All species	Long-tailed Duck	Velvet Scoter
Early autumn	-0.76	-0.61	-0.21
Late autumn	-0.76	-0.60	-0.50
Winter	0.07	-0.03	0.20
Early spring	-0.13	0.16	-0.58
Late spring	-0.53	-0.43	-0.21




Fig. 4. Distribution of all birds observed in autumn 2004 – period average. Explanations as for Fig. 3.

Subsequent controls show pronounced variability of distribution of birds, which is clearly visible in Figure 3, where densities are presented as isolines of different densities – from very low (less than 10 ind./km²) to large localized concentrations (over 200 ind./km²). This can be explained by both feeding conditions and the social behaviour of the birds studied. Despite these substantial fluctuations, the average picture for a season is much less differentiated (Fig. 4). Comparison of the same season in two subsequent years (Fig. 5) shows pronounced variation between years, and it is difficult to say now whether this is mainly due to actual variation in numbers of birds visiting the area in different years or to an inadequate number of observations done in a period of very dynamic changes in migration intensity.

The total average bird distribution in the area is listed in Table 6 and illustrated in Figure 6. It can be seen here that densities occurring within each part of the area vary according to the season, especially in transects 2-3, sector 2. In spring (both early and




Fig. 5. Distributions of all birds observed in November 2003 and 2004 – period average. Explanations as for Fig. 3.

Table 6
Distribution of all birds in the study area in successive seasons.
Densities in sectors are given as numbers of individuals per km²

Early autumn		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	0.0	3.0	2.2	0.7
	5	0.7	3.0	0.7	0.0
	4	0.0	8.9	0.0	0.7
South	3	16.5	4.4	2.2	4.4
	2	76.3	57.8	17.0	10.4
	1	34.1	70.4	93.3	132.6

Late autumn		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	6.3	1.1	7.8	0.7
	5	48.1	0.4	2.5	34.4
	4	11.1	4.1	23.3	24.1
South	3	28.5	15.2	9.3	37.4
	2	33.3	73.0	36.7	31.5
	1	198.1	160.7	419.3	315.6

Table 6 cont.

Winter		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	11.9	57.0	45.6	103.0
	5	2.2	36.3	86.3	180.4
	4	34.8	71.1	29.3	223.3
South	3	20.7	8.3	30.0	21.5
	2	39.3	48.2	24.1	26.7
	1	31.9	85.9	124.8	36.7

Early spring		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	29.6	10.4	121.9	10.4
	5	7.4	8.5	10.7	158.5
	4	28.6	17.0	13.7	111.1
South	3	57.8	27.8	41.9	80.4
	2	40.0	19.6	29.3	68.5
	1	77.0	53.0	63.7	57.4

Early autumn		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	0.0	4.0	0.0	0.0
	5	4.0	3.0	0.0	2.5
	4	1.5	1.0	0.0	4.4
South	3	3.0	21.7	4.9	3.0
	2	8.4	5.9	12.8	0.0
	1	10.9	7.4	5.9	28.1

late) densities are low, in early autumn and winter they are partly low and partly moderate, and only in late autumn does about 25% of the farm area border on more significant densities. Average seasonal densities of the two most abundant species are listed in the Appendix – Tables I and II. Their distributions are varied (Fig. 7-8). In autumn the Long-tailed Duck can be found in high concentrations close to the shore and more to the east of the study area. In winter and spring the highest concentrations of this species can be found farther from the shore, in the easternmost parts of the study area and far from the wind farm location. The water depth in these parts is about 25 m. The Velvet Scoter occurs in early autumn in higher numbers at distances from the shore, but more to the west. Eastern parts of the area are used by this species rather infrequently. In late autumn a high concentration of this species occurs




Fig. 6. Distributions of all birds observed in the area in successive seasons – period averages. Other explanations as for Fig. 3.

near the shore, a bit more to the west. In winter the situation of this species clearly changed, as moderate clusters of this duck were observed only in the northern and north-eastern parts of the study area. In early spring distribution of this species is similar to that observed in autumn, but the numbers are several times lower, reaching




Fig. 7. Distribution of the two commonest species in autumn – period averages. The wind farm location is shown as a rectangle. Other explanations as for Fig. 6.

less than 25 ind./km². In late spring numbers of both species are very low (not more than 10 ind./km²).




Fig. 8. Distribution of the two commonest species in winter and early spring (late spring not shown due to low numbers of birds) – period averages. Explanations as for Fig. 7.

Estimation of the average distribution of birds for all seasons (Fig. 9) shows two distinct concentrations of birds in the study area, both of which are outside the planned location of the wind farm. One of these is on very shallow waters bordering

on the shore, especially in the central and eastern parts of the area. The other is situated in the eastern part, far from the shore, where the average water depth is about 25 m.

The wind farm was planned in an area with relatively low bird densities (under 50 ind./km², in some places even below 25 ind./km²), and only about 30% of the wind farm territory is located within the area of moderate densities.




Fig. 9. Distribution of all species – weighted average for all seasons (explanation in the text – p. XX) – shown on a real map of the area (Fig. 1). Other explanations as for Fig. 6.

Observations of flying birds during ship cruises

As in the case of observations on surface densities (see Fig. 2), flying birds were observed in highly variable numbers (Fig. 10). Single days with high numbers had a strong influence on average values. For all species together the standard deviation was 34.1 while the average was 64.9 ind./km² (so that the coefficient of variation was as high as 52.5%).

The tables in the Appendix (II and III) contain lists of the bird species observed in flight during observations from the ship. The species composition and number relations are similar to that listed for birds observed on the water surface (Table 7). It must be stressed here that the numbers of birds observed flying in winter are not lower than in the migration seasons. This could mean that they are mainly individuals moving between feeding areas and not actively migrating. If this hypothesis is correct, the study area lies outside the main migration routes of waterfowl between north-eastern parts of the Baltic and wintering grounds in the west. In autumn Herring Gulls were quite frequently observed. They were most numerous in November. However, at least some of them did not migrate but were local birds accompanying fishing ships. The numbers of flying birds observed, especially sea ducks, were low in comparison to those observed on the eastern coasts of the Baltic. For example, on the Estonian coast Kontkanen (1996) observed 1,000 flying birds per hour on average during 11 days of




Fig. 10. Densities of birds observed flying (N/h) in each control – for all species together (TOTAL) and for the two most common species. Averages (m) are shown as a line and a 0.05 confidence interval ($\pm 2 \cdot SE$) as a dotted area.

observations. The distribution of flying birds at different distances from the shore (in successive sectors) is presented in Appendix Tables VI-VIII. The highest number was observed in the sector closest to the shore. Seasonal distributions were very similar (Fig. 11) and the same for all species. In the case of the Velvet Scoter, the numbers observed were especially low, which may confirm the earlier suggestion that the area is outside of the main migration route of waterfowl.

Table 7
Comparison of flights and densities of birds in the area studied

		Density	Flights
		Ind./km ²	Ind./h
Total	Autumn	42.9	54.8
	Winter	54.4	58.9
	Spring	54.3	66.8
<i>C. hyemalis</i>	Autumn	24.4	23.6
	Winter	31.2	32.3
	Spring	34.0	25.9
<i>M. fusca</i>	Autumn	14.2	9.0
	Winter	20.4	22.1
	Spring	17.8	27.3

Coastal observations of flying birds

Summer observations of flying birds performed from the coastal point were complementary to the observations from ships. They covered a coastal strip 2 km wide (Table 8, details in Appendix Table IX).

The average flight intensity observed from the Rozewie post was 22.6 ind./hour ($SD = 17.3$). The highest numbers were observed at the end of July and beginning of August (e.g. on 2nd August 267 flying birds were counted within four hours of observation) and on 20th August (Fig. 12). Once again it must be stressed that the variability of daily counts was very high (e.g. on 16th August not one bird was seen for 3 hours). The migration of waterfowl depends greatly on weather conditions (Alerstam 1979, Camphuysen and Dijk 1983, Meissner 2003), but this was not the reason for the variability of the results, as the weather conditions were quite stable during the entire period of observations.

The numbers of flying birds observed were surprisingly low in comparison with the literature data. In Holland and Finland observed numbers (for single species) ranged from 100 ind./h to 300 per hour and more (Camphuysen and Dijk 1983, Buzun 1998). On the German coast the number of passing waterfowl reaches several thousand per day, and in the north-eastern part of the Baltic Sea more than 100,000 individuals can pass an observation post in one day (Moritz 1983, Nehls and Zöllick 1990, Kontkanen 1996).




Fig. 11. Distribution of birds observed flying in successive sectors (1-6) in different seasons

Table 8
Numbers and percentage shares of bird species passing the observation point
at Rozewie in 2002

	No. of individuals	[%]
Diving benthophages		
Velvet Scoter <i>Melanitta fusca</i>	11	1.2
Common Scoter <i>Melanitta nigra</i>	179	19.0
Common Eider <i>Somateria mollissima</i>	170	18.0
Diving ichthyophages		
Black-throated Diver <i>Gavia arctica</i>	1	0.1
Divers <i>indet. Gavia sp.</i>	1	0.1
Great Crested Grebe <i>Podiceps cristatus</i>	7	0.7
Goosander <i>Mergus merganser</i>	155	16.4
Cormorant <i>Phalacrocorax carbo</i>	364	38.6
Others		
Skuas <i>indet. Stercorarius sp.</i>	3	0.3
Mute Swan <i>Cygnus olor</i>	14	1.5
Mallard <i>Anas platyrhynchos</i>	6	0.6
Tufted Duck <i>Aythya fuligula</i>	30	3.2
Marsh Harrier <i>Circus aeruginosus</i>	2	0.2
Grey Heron <i>Ardea cinerea</i>	1	0.1
Total	944	100




Fig. 12. Numbers of birds (N/h) observed during summer observations from the observation point at Jastrzebia Gora

The spatial distribution of flying birds is presented in Table 9. Most of the birds (95%) were observed near the beach. In the farthest strip the average movement was only 1 ind./h. In this strip the most common were sea ducks, and it cannot be ruled out that

they occurred at distances of more than 2 km from the coast. Comparison with the results obtained during ship cruises, however, suggests that the number of birds flying there is still very low. It is possible that the Polish coast is not visited frequently by migrating ducks which can migrate directly from high-concentration localities in Finland, Estonia and Latvia to moulting grounds in the south-western Baltic (Fig. 13).

Table 9
Number of waterfowl flying at different distances from the shore
at the Rozewie observation point

	to 0.5 km	0.5 - 1 km	1 – ca 2 km	Total
Diving benthophages				
Velvet Scoter <i>Melanitta fusca</i>	4	6	1	11
Common Scoter <i>Melanitta nigra</i>	40	132	7	179
Common Eider <i>Somateria mollissima</i>	34	123	13	170
Diving ichthyophages				
Black-throated Diver <i>Gavia arctica</i>			1	1
Divers indet. <i>Gavia</i> sp.			1	1
Great Crested Grebe <i>Podiceps cristatus</i>		5	2	7
Goosanger <i>Mergus merganser</i>	67	86	2	155
Cormorant <i>Phalacrocorax carbo</i>	261	100	3	364
Others				
Skuas indet. <i>Stercorarius</i> sp.		2	1	3
Mute Swan <i>Cygnus olor</i>		6	8	14
Mallard <i>Anas platyrhynchos</i>	6			6
Tufted Duck <i>Aythya fuligula</i>	30			30
Marsh Harrier <i>Circus aeruginosus</i>	1	1		2
Grey Heron <i>Ardea cinerea</i>	1			1
Total	444	461	39	944
[%]	47.0	48.8	4.1	100.0
Individuals/hour	10.6	11.0	1.0	22.6

The passage of birds was observed mainly in the morning and at midday. In these periods 94% of flying birds were observed (Table 10). In the afternoon the most common were sea ducks, but in very low numbers (avg. 3.3 ind./h). Summer migration flights of the Common Scoter on the eastern part of the German coast occur mainly in the afternoon (Nehls and Zöllick 1990). Assuming that the migration speed of this species is around 80 km/h, these birds should pass Rozewie shortly after noon, but this was not the case in our observations. This is another argument supporting the hypothesis about the migration pattern of waterfowl over the Baltic.




Fig. 13. Hypothetical migration flyways of waterfowl in the region

Table 10
Number of waterfowl passing the Rozewie observation point at different times of day

	Morning	Midday	Afternoon	Total
Benthophages				
Velvet Scoter <i>Melanitta fusca</i>	7	3	1	11
Common Scoter <i>Melanitta nigra</i>	92	58	29	179
Common Eider <i>Somateria mollissima</i>	47	112	11	170
Ichthyophages				
Black-throated Diver <i>Gavia arctica</i>	1			1
Divers <i>indet. Gavia sp.</i>	1			1
Great Crested Grebe <i>Podiceps cristatus</i>	4	3		7
Goosanger <i>Mergus merganser</i>	22	133		155
Cormorant <i>Phalacrocorax carbo</i>	245	110	9	364
Others				
Skuas <i>indet. Stercorarius sp.</i>		3		3
Mute Swan <i>Cygnus olor</i>		14		14
Mallard <i>Anas platyrhynchos</i>			6	6
Tufted Duck <i>Aythya fuligula</i>	30			30
Marsh Harrier <i>Circus aeruginosus</i>	1		1	2
Grey Heron <i>Ardea cinerea</i>	1			1
Total	451	436	57	944
[%]	47.8	46.2	6.0	100.0
Individuals/hour	29.6	31.1	4.6	22.6

DISCUSSION

The study area as part of the Baltic Sea winter-quarters of waterfowl

Before evaluating a defined location from the point of view of its suitability for wind farm investments, first the larger area surrounding it should be studied with respect to the migration and wintering system, and only later should local peculiarities be considered. Reliable evaluation of different parts of the Baltic Sea is difficult because of the very different coverage of the area by investigations. For example, Danish and German waters have been thoroughly studied using coast observations, ship surveys and aerial monitoring, while some other areas have been under-studied. Among waters with the poorest coverage is the central Polish coast, where our study area was located. The only available prior data come from a single ship cruise and are undoubtedly imprecise and unreliable. This wholly insufficient data set was the basis for estimations made by Durinck *et al.* (1994) and given in *NATURA 2000* (PLB990002) project documentation. Estimations in these sources are sometimes highly varied, e.g. Durinck *et al.* (*op. cit.*) estimated density of the Velvet Scoter at 42.9 ind./km², while the *NATURA 2000* estimate was 12.5 ind./km², i.e. four times less. The average density of this species within the study area was 23.0 ind./km². For another species, the Long-tailed Duck, estimations by these authors differ by only 10%, but the density of the species in the present study is four times lower. These differences may be due to an insufficient basis for estimations, different areas taken under consideration, and local variation that cannot be shown in large-scale estimations. This strongly underscores the absolute necessity of intensive evaluation of the entire area that can be designated for use for wind energy projects. Only detailed research can indicate which areas are safe enough for birds when used for investments, and such studies are in fact being conducted.

In the general classification by Durinck *et al.* (*op. cit.*), the Polish Baltic coast was ranked 12th among 39 listed, within a numerous group (ranks 6-16) having moderately high scores (15.57 pts., compared to 101.23 pts. for the 5th rank). At the same time, another Polish maritime area (the Pomeranian Bay) is ranked second.

Despite problems with reliable comparisons, published numbers seem to be over-estimated. However, it cannot be ruled out that the area studied was an especially 'empty' location; within this area the average density is only 30.9% of the values given in the *NATURA 2000* project (Table 11). Many observations suggest that the central and eastern parts of the Polish Baltic coast are missed by waterfowl migrating from the north-eastern parts of the Baltic to western and south-western parts of the sea (Fig. 13). However, this is still a hypothesis that should be tested by more extensive and detailed studies.

Table 11
Comparison of the wind farm area with important bird areas on the Baltic Sea

	Area km ²	Total	Ind./km ²	<i>C. hyemalis</i>	Ind./km ²	<i>M. fusca</i>	Ind./km ²
Hoburgs Bank*	1 645	925 770	562.8	925 120	562.4	–	–
Pomeranian Bay*	8 800	1 722 150	195.7	803 000	91.3	357 210	40.6
Kiel Bay*	3 130	449 540	143.6	90 000	28.8	–	–
Szczecin*	3 000	411 140	137.0	–	–	41 300	13.8
Gulf of Riga*	12 000	1 503 440	125.3	1 095 000	91.3	336 000	28.0
Northern Kattegat*	10 500	1 242 120	118.3	–	0.0	82 000	7.8
Gotland*	2 575	262 400	101.9	236 500	91.8	–	–
Polish coast*					89.9		49.2
Natura 2000**	2 116	247 000	116.7	217 000	102.6	26 500	12.5
STUDY AREA	16.2		52.0		29.1		23.0
FARM AREA	4.5		36.1		12.6		21.3
in relation to NATURA	0.21%		30.9%				

* after Durinck *et al.* 1994; ** after NATURA 2000 PLB990002

Local distribution patterns

In addition to the characteristics of the larger region, it is essential to know the local peculiarities of the distribution of birds. Even in a region generally important for birds, local conditions (e.g. water depth, geology, or feeding conditions) may make a particular location designated for a wind farm sufficiently safe for the bird population. Conversely, an area that is generally not of great importance for birds may contain special sites where high concentrations of birds can be found. Such situations cannot be predicted with high probability without first conducting special research at the site. The present evaluation is an excellent example, as within the study area there are places where bird concentrations are quite high and others with permanently low densities. Similar were conclusions by Meissner (2010), although documented using different presentation method.

The species composition found in the area is consistent with accidental observations done here at the end of the 1960s and beginning of the 1970s and again at the end of the 1980s and beginning of the 1990s, and is associated with patterns in feeding conditions. More than 93% of the birds observed were of only two species: Long-tailed Duck and Velvet Scoter. The Common Scoter and Herring Gull had shares of more than 1%. The first of these species was associated with the shallowest waters, while Herring Gulls were associated with fishing activity.

The average densities in the study area can be described as moderate. The highest concentrations of waterfowl were near the coast and to the east of the planned wind farm location. However, sometimes larger flocks of Velvet Scoter visited the area.

Comparison of the same season in two subsequent years shows a considerable fluctuation in densities from year to year, but it is still unknown whether this reflects actual changes in numbers (e.g. caused by feeding conditions – Kube and Skov 1996) or weak-

ness of the sampling, i.e. too few cruises in relation to the very dynamic process of bird migration. This aspect is still not well understood and requires further study.

The results of the observations show that the waterfowl migration is not too intensive here, especially in comparison with the data from the eastern Baltic. All gulls and terns migrate along the seashore. Here too intensive migration of waterfowl was not very intensive – it seems that they miss the area and shorten their way from the north-eastern Baltic to western and south-western parts of the sea by flying directly over the open sea. According to Kontkanen (1996), the Long-tailed Duck can migrate during the night, but such migration should be accompanied by diurnal migration as well. This was not found during the observations, at least in the area studied.

The relatively small differences in the numbers of birds observed in flight in winter and during migration periods suggest that actual migration is weak here and most of the birds seen were making local feeding movements.

The results of the research show that the location studied is relatively unattractive to waterfowl (about 31% of the average density estimated in the *NATURA 2000* project) and lies adjacent to areas that are more attractive (around 70% of the average), but still not very important for birds.

REFERENCES

- Alerstam T. 1979. *Wind as a selective agent in bird migration*. *Ornis Scand.* 10: 76-93.
- Buzun V. A. 1998. *Autumn migration of birds over the Koorgalsky Reef Archipelago (south-eastern Gulf of Finland) in 1997*. Proc. of the Programme: "Study of the status & Trends of Migratory Bird Populations in Rusia", Moscow. pp. 108-121.
- Camphuysen C. J., Dijk J. 1983. *Zee- en kustvogels langs de Nederlandste kust, 1974-79*. *Limosa* 56: 81-230.
- Durinck J., Skov H., Jensen F. P., Pihl S. 1994. *Important marine areas for wintering birds in the Baltic Sea*. *Ornis Consult.* Copenhagen.
- Heinemann D. 1981. *A range finder for pelagic bird censusing*. *Journal of Wildlife Management* 45: 489-493.
- Komdeur J., Bertelsen J., Cracknell G. 1992. *Manual for Aeroplane and Ship Surveys of Waterfowl and Seabirds*. IWRB Special Publication No. 19.
- Kontkanen H. 1996. *Autumn migration of Arctic waterfowl in Finland, Estonia and northwestern Russia in 1993*. *Wetlands International Seaduck Specialist Group Bull.* 6: 32-41.
- Kube J., Skov H. 1996. *Habitat selection, feeding characteristics, and food consumption of long-tailed ducks, Clangula hyemalis, in the southern Baltic Sea*. *Meereswissenschaftliche Berichte* 18: 83-100.
- Meissner W. 2003. *Wiosenny przelot mew Laridae koło przylądka Rozewie*. *Not. Orn.* 44: 179-186.
- Meissner W. 2010. *Seasonal changes in numbers and distribution of the Long-tailed Duck Clangula hyemalis, Common Scoter Melanitta nigra and Velvet Scoter M. fusca near the Cape Rozewie*. *Ornis Polonica* 51: 275-284.
- Moritz D. 1983. *Vom Mauserzug der Eiderente (Somateria mollissima) an der Ostseeküste Schleswig-Holsteins bei Schleimünde*. *Seevögel* 4: 57-64.
- Nehls H. W., Zöllick H. 1990. *The moult migration of the Common Scoter (Melanitta nigra) off the coast of the GDR*. *Proc. Conf. Baltic Birds* 5. pp. 36-46. Riga.

APPENDIX Table I

Distribution of Long-tailed Ducks in the study area in successive seasons.

Densities in sectors are given as numbers of individuals per km²

Early autumn		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	0.0	3.0	0.0	0.0
	5	0.7	1.5	0.0	0.0
	4	0.0	0.7	0.0	0.0
	3	0.0	0.0	0.0	3.7
South	2	0.0	0.0	1.5	2.2
	1	0.0	59.3	75.6	117.0
Late autumn		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	0.0	0.0	1.5	0.4
	5	1.1	0.0	0.4	27.0
	4	0.0	0.4	1.1	8.5
	3	0.0	0.0	0.0	16.3
South	2	0.0	0.4	1.9	5.9
	1	1.1	125.2	412.2	305.6
Winter		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	10.4	6.9	26.7	95.9
	5	0.0	0.0	5.2	142.2
	4	0.0	11.1	12.6	148.1
	3	0.0	2.2	5.6	19.3
South	2	4.4	20.0	5.2	20.4
	1	0.0	64.4	89.3	24.8
Early spring		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	27.2	4.1	120.4	9.3
	5	4.9	0.4	4.1	155.2
	4	5.4	8.1	7.4	104.4
	3	6.9	13.0	14.8	73.3
South	2	2.0	3.7	7.0	64.8
	1	0.0	8.5	18.1	41.9

Table II cont.

Late spring		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	0.0	0.0	0.0	0.0
	5	0.0	0.0	0.0	1.0
	4	0.0	0.0	0.0	4.4
South	3	1.0	1.0	1.0	3.0
	2	2.0	0.0	2.0	0.0
	1	5.9	1.0	1.0	26.2

APPENDIX Table II

Distribution of Velvet Scoters in the study area in successive seasons.

Densities in sectors are given as numbers of individuals per km²

Early autumn		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	0.0	0.0	2.2	0.0
	5	0.0	1.5	15.6	0.0
	4	0.0	5.2	0.0	0.0
South	3	16.3	3.7	0.0	0.0
	2	71.1	56.3	0.0	0.0
	1	14.1	7.4	0.0	0.0
Late autumn		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	3.7	0.0	0.0	0.0
	5	0.7	0.0	1.5	5.2
	4	7.8	2.6	21.1	11.9
South	3	28.5	11.5	9.3	19.6
	2	31.1	67.8	32.6	23.3
	1	175.6	25.2	3.0	8.1
Winter		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	1.5	51.1	18.1	5.9
	5	0.0	32.6	75.6	9.3
	4	34.1	57.0	5.9	73.3
South	3	20.7	4.4	23.7	1.1
	2	34.1	23.7	18.9	0.7
	1	28.9	19.3	34.8	7.4

Table II cont.

Early spring		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	2.5	6.3	1.5	0.7
	5	2.5	7.8	6.7	3.3
	4	23.2	8.5	6.3	5.9
	3	49.9	14.8	26.7	6.7
South	2	38.0	15.9	21.9	3.0
	1	42.5	31.9	43.3	14.1
Early autumn		Transects			
		West		East	
North	Sectors	1 18°00'	2 18°06'	3 18°12'	4 18°18'
	6	0.0	4.0	0.0	0.0
	5	4.0	3.0	0.0	0.0
	4	0.0	1.0	0.0	0.0
	3	2.0	20.7	4.0	0.0
South	2	6.4	5.9	10.9	0.0
	1	3.5	4.4	3.0	0.0

APPENDIX Table III
Species composition and numbers of birds observed flying in the Dębki area
in autumn of 2003 and 2004

	October		November				Total
	4 Oct. 2004	25 Oct. 2004	4 Nov. 2004	14 Nov. 2003	28 Nov. 2003	29 Nov. 2004	
Diving benthophages							
Long-tailed Duck <i>Clangula hyemalis</i>		16	278	66	124	83	567
Velvet Scoter <i>Melanitta fusca</i>	2	73	68	34	11	29	217
Common Scoter <i>Melanitta nigra</i>	8	13		2			23
Common Eider <i>Somateria mollissima</i>			2	151	2	2	157
Diving ichthyophages							
Black-throated Diver <i>Gavia arctica</i>	1	1	1				3
Red-throated Diver <i>Gavia stellata</i>				3		1	4
Divers <i>indet. Gavia sp.</i>		2	5	1	4	2	14
Goosanger <i>Mergus merganser</i>					25		25
Red-breasted Merganser <i>Mergus serrator</i>				2	4		6
Cormorant <i>Phalacrocorax carbo</i>	2						2
Others							
Herring Gull <i>Larus argentatus</i>	19	36	77	46	82	50	310
Common Gull <i>Larus canus</i>	1	2	1	4	3	2	13
Great Black-backed Gull <i>Larus marinus</i>		2	2	2	2	1	9
Little Gull <i>Larus minutus</i>		1					1
Kittiwake <i>Rissa tridactyla</i>				1			1
Black Tern <i>Chlidonias niger</i>	1						1
Swans <i>indet. Cygnus sp.</i>				3			3
Widgeon <i>Anas penelope</i>		30					30
Mallard <i>Anas platyrhynchos</i>	15						15
Tufted Duck <i>Aythya fuligula</i>			1				1
Total	49	176	435	315	257	170	1402
Individuals/hour	12.3	44.0	108.8	78.8	64.3	42.5	58.4

APPENDIX Table IV
Species composition and numbers of birds observed flying in the Dębki area
in winter 2003/2004

	December		January	February	Total
	9 Dec. 2003	30 Dec. 2003	27 Jan. 2004	27 Feb. 2004	
Diving benthophages					
Long-tailed Duck <i>Clangula hyemalis</i>	45	93	190	124	452
Velvet Scoter <i>Melanitta fusca</i>	49	87	72	102	310
Common Scoter <i>Melanitta nigra</i>	12	3	5	8	28
Common Eider <i>Somateria mollissima</i>				17	17
Diving ichthyophages					
Black-throated Diver <i>Gavia stellata</i>	2	1			3
Divers <i>indet. Gavia sp.</i>		1	4		5
Cormorant <i>Phalacrocorax carbo</i>				1	1
Others					
Great Black-backed Gull <i>Larus marinus</i>		2			2
Herring Gull <i>Larus argentatus</i>	19	38	34	17	108
Common Gull <i>Larus canus</i>	2	2	4		8
Whooper Swan <i>Cygnus cygnus</i>		6			6
Swans <i>indet. Cygnus sp.</i>		2			2
Total	129	235	309	269	942
Individuals/hour	64.5	58.8	77.3	67.3	58.9

APPENDIX Table V
Species composition and numbers of birds observed flying in the Dębki area
in spring 2004

	5 March	28 March	5 April	20 April	5 May	Total
Diving benthophages						
Long-tailed Duck <i>Clangula hyemalis</i>	60	195	180	46	12	493
Velvet Scoter <i>Melanitta fusca</i>	18	220	176	59	45	518
Common Scoter <i>Melanitta nigra</i>		2	3	5	80	90
Diving ichthyophages						
Black-throated Diver <i>Gavia arctica</i>	1		3	5	3	12
Red-throated Diver <i>Gavia stellata</i>			3			3
Divers <i>indet.</i> <i>Gavia sp.</i>	2	1	7		8	18
Cormorant <i>Phalacrocorax carbo</i>					1	1
Others						
Great Black-backed Gull <i>Larus marinus</i>		1				1
Herring Gull <i>Larus argentatus</i>	9	25	20	14	14	82
Common Gull <i>Larus canus</i>	1		11			12
Black-headed Gull <i>Larus ridibundus</i>			1			1
Little Gull <i>Larus minutus</i>			6	1	18	25
Kittiwake <i>Rissa tridactyla</i>					2	2
Black Tern <i>Chlidonias niger</i>					11	11
Total	91	444	410	130	194	1269
Ind./hour	22.8	148.0	102.5	32.5	48.5	66.8

APPENDIX Table VI
Species composition and numbers of birds observed flying in the Dębki area
in winter within subsequent sectors

Sectors:	Distance from the shore [km]						Total
	1.5-3.8	3.8-5.1	5.1-7.4	7.4-9.7	9.7-12.0	12.0-14.3	
	1	2	3	4	5	6	
Diving benthophages							
Long-tailed Duck <i>Clangula hyemalis</i>	165	101	98	52	15	21	452
Velvet Scoter <i>Melanitta fusca</i>	62	82	75	66	20	5	310
Common Scoter <i>Melanitta nigra</i>	28						28
Common Eider <i>Somateria mollissima</i>	7	10					17
Diving ichthyophages							
Black-throated Diver <i>Gavia stellata</i>		1		2			3
Divers <i>indet. Gavia sp.</i>	1	1	3				5
Cormorant <i>Phalacrocorax carbo</i>	1						1
Others							
Great Black-backed Gull <i>Larus marinus</i>	1		1				2
Herring Gull <i>Larus argentatus</i>	21	12	18	19	26	12	108
Common Gull <i>Larus canus</i>		1	2	1	4		8
Whooper Swan <i>Cygnus cygnus</i>	6						6
Swans <i>indet. Cygnus sp</i>	2						2
Total	294	208	197	140	65	38	942
Individuals/hour	108.9	77.0	73.0	51.9	24.1	14.1	58.9

APPENDIX Table VII
Species composition and numbers of birds observed flying in the Dębki area
in autumn within successive sectors

Sectors:	Distance from the shore [km]						Total
	1.5-3.8	3.8-5.1	5.1-7.4	7.4-9.7	9.7-12.0	12.0-14.3	
	1	2	3	4	5	6	
Diving benthophages							
Long-tailed Duck <i>Clangula hyemalis</i>	185	96	90	57	84	55	567
Velvet Scoter <i>Melanitta fusca</i>	22	58	65	41	18	13	217
Common Scoter <i>Melanitta nigra</i>	15	3				5	23
Common Eider <i>Somateria mollissima</i>	155	2					157
Diving ichthyophages							
Black-breasted Diver <i>Gavia arctica</i>		3					3
Red-breasted Diver <i>Gavia stellata</i>		4					4
Divers <i>indet. Gavia sp.</i>	1		2	5	1	5	14
Goosanger <i>Mergus merganser</i>						25	25
Red-breasted Merganser <i>Mergus serrator</i>		5	1				6
Cormorant <i>Phalacrocorax carbo</i>		1				1	2
Others							
Great Black-backed Gull <i>Larus marinus</i>		3	2	1	3		9
Herring Gull <i>Larus argentatus</i>	54	47	57	58	61	33	310
Common Gull <i>Larus canus</i>	3	2	1	2	4	1	13
Little Gull <i>Larus minutus</i>						1	1
Kittiwake <i>Rissa tridactyla</i>		1					1
Black Tern <i>Chlidonias niger</i>		1					1
Swans <i>indet. Cygnus sp.</i>			3				3
Widgeon <i>Anas penelope</i>				30			30
Mallard <i>Anas platyrhynchos</i>					3	12	15
Tufted Duck <i>Aythya fuligula</i>		1					1
Total	435	227	221	194	174	151	1402
Individuals/hour	108.8	56.8	55.3	48.5	43.5	37.8	58.4

APPENDIX Table VIII
Species composition and numbers of birds observed flying in the Dębki area
in spring within subsequent sectors

Sectors:	Distance from the shore [km]						Total
	1.5-3.8	3.8-5.1	5.1-7.4	7.4-9.7	9.7-12.0	12.0-14.3	
	1	2	3	4	5	6	
Diving benthophages							
Long-tailed Duck <i>Clangula hyemalis</i>	203	63	109	42	33	43	493
Velvet Scoter <i>Melanitta fusca</i>	150	88	92	112	60	16	518
Common Scoter <i>Melanitta nigra</i>	55	35					90
Diving ichthyophages							
Black-breasted Diver <i>Gavia arctica</i>	8		1	1		2	12
Red-breasted Diver <i>Gavia stellata</i>	1		1	1			3
Divers <i>indet. Gavia sp.</i>		9	3	3	2	1	18
Cormorant <i>Phalacrocorax carbo</i>	1						1
Others							
Great Black-backed Gull <i>Larus marinus</i>						1	1
Herring Gull <i>Larus argentatus</i>	23	16	15	10	6	12	82
Common Gull <i>Larus canus</i>	1	3	4	1	3		12
Black-headed Gull <i>Larus ridibundus</i>				1			1
Little Gull <i>Larus minutus</i>			2	3	14	6	25
Kittiwake <i>Rissa tridactyla</i>					2		2
Black Tern <i>Chlidonias niger</i>	11						11
Total	453	214	227	174	120	81	1269
Individuals/hour	142.9	67.5	71.6	54.9	37.9	25.6	66.8

APPENDIX Table IX
Details of observations made in summer 2002 at the Rozewie observation point

	27 Jul.	30 Jul.	2 Aug.	5 Aug.	9 Aug.	12 Aug.	16 Aug.	20 Aug.	24 Aug.	27 Aug.	31 Aug.	4 Sep.	Total
Benthophages													
Velvet Scoter <i>Melanitta fusca</i>	2			2	1						1	5	11
Common Scoter <i>Melanitta nigra</i>	15	7	36	23		11		25	12		1	49	179
Common Eider <i>Somateria mollissima</i>		75	51	3	20	6		1	12	2			170
Ichthyophages													
Black-throated Diver <i>Gavia arctica</i>									1				1
Divers indet. <i>Gavia</i> sp.						1							1
Great Crested Grebe <i>Podiceps cristatus</i>	2		3	2									7
Goosander <i>Mergus merganser</i>	13	11	90		5	6		1	6	4	19		155
Cormorant <i>Phalacrocorax carbo</i>	114	14	86	6	16	21		67	15	7	11	7	364
Others													
Skuas indet. <i>Stercorarius</i> sp.										3			3
Mute Swan <i>Cygnus olor</i>	14												14
Mallard <i>Anas platyrhynchos</i>						6							6
Tufted Duck <i>Aythya fuligula</i>	16			11	3								30
Marsh Harrier <i>Circus aeruginosus</i>								1				1	2
Grey Heron <i>Ardea cinerea</i>			1										1
Total	176	107	267	47	45	51	0	95	46	16	32	62	944
Number of observation hours per day	5:30	4:15	4:00	3:30	3:00	3:00	3:00	3:00	3:00	3:00	3:00	3:30	41:45
Individuals/hour	32.0	25.2	66.8	13.4	15.0	17.0	0.0	31.7	15.3	5.3	10.7	17.7	22.6